

Distinct Scaling Regimes of Energy Release Dynamics in the Nighttime Magnetosphere

V. M. Uritsky¹, E. Donovan¹, A. J. Klimas², and E. Spanswick¹

Based on a spatiotemporal analysis of POLAR UVI images, we show that the auroral emission events that initiate equatorward of the isotropic boundary (IB) obtained from a time-dependent empirical model, have systematically steeper power-law slopes of energy, power, area and lifetime probability distributions compared to the events that initiate poleward of the IB. The low-latitude group of events contains a distinct subpopulation of substorm-scale disturbances violating the power-law behavior, while the high latitude group is described by nearly perfect power-law statistics over the entire range of scales studied. The results obtained indicate that the inner and outer portions of the plasma sheet are characterized by substantially different scaling regimes of bursty energy dissipation suggestive of different physics in these regions.

1. Introduction

The activity of the nighttime auroral oval represents a wide range of dynamical processes in the magnetotail, including substorm expansion onsets, pseudobreakups, steady magnetospheric convection events with or without substorms, bursty bulk flows, and sawtooth events (see e.g., *Zesta et al.* [2000]; *Lui* [2001]; *Frey et al.* [2004]; *Henderson et al.* [2006]). Despite the diversity of physical conditions associated with each particular type of auroral activity, their net energy output can be described by a set of apparently universal power-laws (*Lui et al.* [2000]; *Lui* [2002]; *Uritsky et al.* [2003, 2002, 2006]) signaling the existence of an organizing dynamical principle arranging intermittent magnetospheric dissipation across vast ranges of spatial and temporal scales.

Power-law intermittency of energy dissipation has attracted significant attention in modern statistical mechanics (see *Dhar* [2006] and refs therein) and is often considered a hallmark of turbulent and/or critical phenomena with no characteristic scales other than those dictated by the finite size of the system (*Sreenivasan et al.* [2004]; *Lubeck* [2004]). Examples of such behavior in geo- and space sciences include fully developed turbulence in hydrodynamic or magnetized flows (*Lazarian* [2006]), Gutenberg-Richter statistics of earthquake magnitudes (*Turcotte* [1989]), scale-invariance in the solar corona (*Charbonneau et al.* [2001]). In this context, the auroral activity provides one of the most impressive

¹Physics and Astronomy Department, University of Calgary, Calgary, AB, Canada

²UMBC at NASA / Goddard Space Flight Center, Greenbelt, Maryland, USA

examples of scale-free behavior in nature. The energy distribution of electron emission regions exhibits a power-law shape over a range of 6 orders of magnitude (*Uritsky et al.* [2002]) which can be extended to up to 11 orders by combining the satellite data with ground-based TV observations (*Kozelov et al.* [2004]).

The auroral emission statistics reported so far represent global long-term properties of nighttime magnetospheric disturbances. The fact that these properties are dominated by power-law scaling does not eliminate the possibility of a more complex behavior on the level of specific plasma sheet structures described by drastically different physical conditions and geometry. In this study, we are taking a step toward a better understanding of the relationship between the scale-free auroral precipitation statistics and the underlying central plasma sheet (CPS) morphology. We suggest that the inner and the outer CPS regions are responsible for three distinct scaling modes of the auroral precipitation dynamics, and provide a possible physical interpretation for the observed differences.

2. Data and Algorithm

We have studied time series of digital images of nighttime northern aurora (55-80 MLat, 2000 - 0400 MLT) taken by the Ultraviolet Imager (UVI) onboard the POLAR spacecraft in the 165.5 to 174.5 nm portion of the Lyman-Birge-Hopfield spectral band (integration time 36.5 s, time resolution 184 s). The data analyzed include 16,000 images covering two observation periods: 01/01/1997 - 02/28/1997 and 01/01/1998 - 02/28/1998. Our analysis was based on spatiotemporal tracking of auroral emission events (*Uritsky et al.* [2002, 2003]). The UV luminosity $w(t, \mathbf{r})$ was studied as a function of time t and position $\mathbf{r}$ on the image plane. First, active auroral regions were identified by applying an activity threshold w_a representing a background UV flux. Contiguous spatial regions with $w(\mathbf{r}, t) > w_a$ were treated as pieces of evolving events. Second, by checking for overlap of common pixels between each pair of consecutive UVI frames, we constructed a set of spatiotemporal integration domains $\Lambda_i (i = 1, \dots, N)$ corresponding to each of N individual emission events found by our method. These domains of contiguous activity in space and time were used to compute the lifetime, T_i , the energy, $E_i = k \int_{\Lambda_i} w(\mathbf{r}, t) d\mathbf{r} dt$, the peak power, $W_i = k \max_t (\int_{\Lambda_i(t)} w(\mathbf{r}, t) d\mathbf{r})$, as well the peak area, $A_i = \max_t (\int_{\Lambda_i(t)} d\mathbf{r})$ of every event, where $k = 2.74 \times 10^{-8} \text{ J} \cdot \text{photon}^{-1}$ (*Brittnacher et al.* [1997]) and the integrals were numerically approximated by sums. The statistics reported below are for the threshold $w_a = 10 \text{ photons} \cdot \text{cm}^2 \cdot \text{s}^{-1}$. Their main features remain the same if the threshold is varied at least within the range 5 to 15 $\text{photons} \cdot \text{cm}^2 \cdot \text{s}^{-1}$.

The auroral onset positions of each event were estimated with an error of about 300 km in either spatial direction. We organized the data to allow us to compare events that likely originate in the inner magnetosphere relative to events that likely originate in the outer magnetosphere. To accomplish this, we determine the event locations relative to the isotropic boundary (IB) at the meridian of the event origin ¹:

$$\phi_i = MLat_i - [A_0 - A_1 \cos(\pi(MLT_i - MLT_0)/12)] \quad (1)$$

Here, MLT_i and $MLat_i$ are the onset coordinates of the

i -th event. A_0 , A_1 , MLT_0 are the coefficients of the empirical IB model due to *Gvozdevsky and Sergeev* [1995] which were computed based on hourly values of auroral electrojet index (WDC for Geomagnetism, Kyoto) and solar wind dynamic pressure (ACE spacecraft) at the beginning of each auroral intensification. The ions maintain their isotropy on the poleward side of IB due to the effective pitch angle scattering in the tail current sheet (*Newell et al.* [1996]). As a result, the latitude of IB has a high correlation with the magnetic field inclination at the geomagnetic equator as well as with the equatorward boundary of the proton aurora as determined from direct optical observations (*Donovan et al.* [2003]). Here, we have used the empirical IB model to separate inner and outer emission events. Positive values of ϕ on average indicate events originating on field lines that map further out on the plasma sheet, as opposed to negative values corresponding to the events that originate closer to Earth.

The statistics of the emission events that initiated on the poleward ($\phi > 0$) and on the equatorward ($\phi < 0$) side of the IB are characterized by sets of probability density distributions $p(x)$, where $x \in \{E, W, A, T\}$. The power-law exponents obtained from these distributions are denoted as τ_x , with the subscript indicating the variable under study.

3. Results and Discussion

In the end we have 7481 events, 6231 and 1250 poleward and equatorward of the IB, respectively. These yielded a database of parameters E , W , A , T , MLT , $MLat$, ϕ for those events. Figure 1 is the emission energy E as a function of the relative onset latitude ϕ for all events ($n = 7481$). The scatterplot has clear asymmetric shape suggesting different statistics for negative and positive ϕ values. As we discuss below, these statistical subsets are characterized by significantly different regimes of scaling behavior indicating different physical environments.

Figure 2 shows probability distributions for the events which initiated on the poleward and equatorward sides of the IB. The high-latitude (HL) events ($\phi > 0$) exhibit stable power-law distributions of emission energy E , peak emission power W , lifetime T and peak area A (not shown) with near-zero regression errors. The ranges of scales of these power-law behaviors involves both small auroral activations and large events whose energy output lies in the range of global substorms. The exponents τ_E , τ_W , τ_A and τ_T are close to the values obtained earlier for the same observation period without filtering the activity by the onset location (*Uritsky et al.* [2002]).

The low-latitude (LL) events ($\phi < 0$) have more complicated statistical properties as can be expected from Fig. 1. Their distribution functions demonstrate a crossover behavior involving small-scale regions described by τ_x exponents which are greater than the corresponding exponents of HL events, as well as large-scale regions where the slopes are significantly shallower (the τ_x exponents for these regions are not well defined due to the insufficient number of large-scale LL events). The position of the crossover $E^* = 5 \times 10^{12}$ J on the $p(E)$ distribution of LL events matches the data gap in Fig. 1 separating low- and high-energy events with negative relative latitudes ϕ .

As can be seen from Table 1, only 1/6 of the events contribute to the population with $\phi < 0$. However, their

participation in the energy budget of the electron aurora approaches 50 percent as can be seen from the comparison of the overall energy E_{tot} released by LL and HL events. The fact that the HL events are the most common form of emission dynamics (relative occurrence frequency 82.3%) is not a surprise as these events likely map to the reconnection regions of the plasma sheet, as we briefly discuss below. The table also shows that the scaling behaviors of LL and LH events are described by substantially different sets of τ exponents.

Table 2 provides scaling exponents of energy distributions $p(E)$ as well as several parameters reflecting the state of the solar wind and the auroral magnetosphere at the beginnings of emission events for four non-overlapping ranges of magnetic latitude $MLat$. The data show that the group of events that initiated southward of $MLat=65$ are characterized by significantly higher statistical values of solar wind dynamic pressure and electrojet index. These events seem to appear in a rather stretched magnetotail configuration as reflected by the magnetotail (MT) index (*Gvozdevsky and Sergeev [1995]*). One can also notice that the lower the magnetic latitude, the higher is the exponent τ_E . This tendency is in accordance with the difference in the LL- and HL-event statistics reported in Table 1, and it also suggests that the change of the scaling regime of auroral emission events with onset latitude is a continuous process rather than a discrete transition fully dictated by the empirical IB location.

From the statistical mechanics viewpoint, the existence of two or more scaling regimes within the same physical system signals the presence of several distinct universality classes (UC) in the underlying turbulent dynamics (*Lubeck [2004]; Dhar [2006]*). It is known that many stochastic scale-invariant phenomena can be mapped onto a finite collection of such classes. This mapping is insensitive to a variety of physical parameters and conditions and can be the same for quite different systems. Typically, each UC brings its own set of critical exponents which express its inherent set of relevant variables and the underlying small- and large-scale symmetries (*BenHur and Biham [1996]*). In this context, our results clearly show that the energy release dynamics in the inner and outer CPS belong to different UCs and are therefore governed by different physical mechanisms.

The nature of the isotropic boundary sheds some light on the scaling regimes observed. On average, HL events tend to initiate in the outer CPS region. The energy conversion in this region is believed to be dominated by magnetic reconnection (*Birn and Hones [1981]*). Our results strongly suggest that near-Earth (midtail) reconnection is a turbulent bursty process with no well-defined dissipation scales. Judging by the relative occurrence of these events (Table 1), the scale-free activity in the reconnection region is the dominant mode of energy release dynamics in the nightside magnetosphere. The power-law emission exponents obtained for this region are rather close to the exponents from a driven current sheet simulation (*Klimas et al. [2004]*) consistent with multiscale turbulent reconnection being the source of scale-invariance in the outer CPS dynamics. The LL events are mainly produced in the inner CPS. With a less stretched magnetic field topology, this region is not a region that favors magnetic reconnection; however, it can be prone to current disruption which offers an alternate mechanism for energy release in the inner tail (*Lui [2000]*).

Our findings can be summarized as follows: (1) the high-latitude emission events are characterized by broadband power-law statistics with no characteristic scales;

(2) the low-latitude events constitute a non-uniform statistical population with an energy crossover separating large and small scale activity; (3) the scaling exponents of the low- and high-latitude events are significantly different within the entire range of scales studied, which signals distinct universality classes of energy release dynamics. To the extent that the empirical IB model can be considered an accurate representation of the true IB, the onsets of the HL events are located in an outer CPS region, the onsets of the LL events in an inner CPS regions, and these two regions exhibit distinctly different energy release dynamics.

The difference in UC following from this analysis is suggestive of different physics. It would be a significant step forward if we could use physics-based models of CPS dynamics to predict UCs, and to thereby determine how the different statistical behaviors of the inner and outer regions might be generated.

Acknowledgments. This work was partly supported by NSERC operating grant of EFD. We thank William Liu for stimulating discussions.

Notes

1. A more accurate analysis based on Tsyanenko T96 and T05 field-line models as well as optical determination of the isotropic boundary will be published elsewhere.

References

- BenHur, A., and O. Biham (1996), Universality in sandpile models, *Phys. Rev. E*, *53*(2), R1317–R1320.
- Birn, J., and E. W. Hones (1981), 3-dimensional computer modeling of dynamic reconnection in the geomagnetic tail, *J. Geophys. Res.*, *86*(NA8), 6802–6808.
- Brittnacher, M., R. Elsen, G. Parks, L. Chen, G. Germany, and J. Spann (1997), A dayside auroral energy deposition case study using the POLAR Ultraviolet imager, *Geophysical Research Lett.*, *24*(8), 991–994.
- Charbonneau, P., S. W. McIntosh, H. L. Liu, and T. J. Bogdan (2001), Avalanche models for solar flares, *Solar Phys.*, *203*(2), 321–353.
- Dhar, D. (2006), Theoretical studies of self-organized criticality, *Physica A*, *369*(1), 29–70.
- Donovan, E. F., B. J. Jackel, I. Voronkov, T. Sotirelis, F. Creutzberg, and N. A. Nicholson (2003), Ground-based optical determination of the b2i boundary: A basis for an optical MT-index, *J. Geophys. Res.*, *108*(A3), doi:10.1029/2001JA00919.
- Frey, H. U., S. B. Mende, V. Angelopoulos, and E. F. Donovan (2004), Substorm onset observations by IMAGE-FUV, *J. Geophys. Res.*, *109*(A10), doi:10.1029/2004JA010607.
- Gvozdevsky, B. B., and V. A. Sergeev (1995), MT-index – A possible new index to characterize the configuration of the magnetotail, *Three-Dimensional Magnetosphere*, *18*(8), 51–54.
- Henderson, M. G., et al. (2006), Substorms during the 10-11 August 2000 sawtooth event, *J. Geophys. Res.*, *111*(A6), doi:10.1029/2005JA011366.
- Klimas, A. J., V. M. Uritsky, D. Vassiliadis, and D. N. Baker (2004), Reconnection and scale-free avalanching in a driven current-sheet model, *J. Geophys. Res.*, *109*(A2), doi:10.1029/2003JA010036.
- Kozelov, B. V., V. M. Uritsky, and A. J. Klimas (2004), Power law probability distributions of multiscale auroral dynamics from ground-based TV observations, *Geophysical Research Lett.*, *31*(20), doi:10.1029/2004GL020962.
- Lazarian, A. (2006), Intermittency of magnetohydrodynamic turbulence: An astrophysical perspective, *Int. J. Modern Phys. D*, *15*(7), 1099–1111.

- Lubeck, S. (2004), Universal scaling behavior of non-equilibrium phase transitions, *Int. J. Modern Phys. B*, *18*(31-32), 3977–4118.
- Lui, A. T. Y. (2000), Electric current approach to magnetospheric dynamics and the distinction between current disruption and magnetic reconnection, in *AGU Monogr. Ser.*, vol. 118, edited by S. Ohtani, R. Fujii, M. Hesse, and R. L. Lysak, pp. 31–40, AGU, Washington, DC.
- Lui, A. T. Y. (2001), Current controversies in magnetospheric physics, *Rev. Geophysics*, *39*(4), 535–563.
- Lui, A. T. Y. (2002), Multiscale phenomena in the near-Earth magnetosphere, *J. Atmospheric Solar - Terrestrial Phys.*, *64*(2), 125–143.
- Lui, A. T. Y., S. C. Chapman, K. Liou, and et al. (2000), Is the dynamic magnetosphere an avalanching system?, *Geophys. Res. Lett.*, *27*, 911–914.
- Newell, P. T., Y. I. Feldstein, Y. I. Galperin, and C. I. Meng (1996), Morphology of nightside precipitation, *J. Geophys. Res.*, *101*(A5), 10,737–10,748.
- Sreenivasan, K. R., A. Bershadskii, and J. J. Niemela (2004), Multiscale SOC in turbulent convection, *Physica A*, *340*(4), 574–579.
- Turcotte, D. L. (1989), Fractals in geology and geophysics, *Pure Appl. Geophysics*, *131*(1-2), 171–196.
- Uritsky, V. M., A. J. Klimas, D. Vassiliadis, D. Chua, and G. Parks (2002), Scale-free statistics of spatiotemporal auroral emissions as depicted by POLAR UVI images: Dynamic magnetosphere is an avalanching system, *J. Geophys. Res.*, *107*(A12), doi:10.1029/2001JA000281.
- Uritsky, V. M., A. J. Klimas, and D. Vassiliadis (2003), Evaluation of spreading critical exponents from the spatiotemporal evolution of emission regions in the nighttime aurora, *Geophys. Res. Lett.*, *30*(15), doi:10.1029/2002GL016556.
- Uritsky, V. M., A. J. Klimas, and D. Vassiliadis (2006), Critical finite-size scaling of energy and lifetime probability distributions of auroral emissions, *Geophys. Res. Lett.*, *33*(8), doi:10.1029/2005GL025330.
- Zesta, E., L. R. Lyons, and E. Donovan (2000), The auroral signature of Earthward flow bursts observed in the magnetotail, *Geophys. Res. Lett.*, *27*(20), 3241–3244.

V. M. Uritsky, Physics and Astronomy Department, University of Calgary, SB605, 2500 University Drive NW, Calgary, AB T3A0P4, Canada. (vuritsky@phas.ucalgary.ca)




Figure 1. Scatterplot of emission energies E versus relative magnetic latitudes ϕ of onset locations measured with respect to the empirical IB latitude (IBL) (see eq.(1)). The high-latitude ($\phi > 0$) and the low-latitude ($\phi < 0$) events contribute to two distinct scaling regimes of energy release dynamics in the nighttime magnetosphere as discussed further in the text. The dashed horizontal line marks the position of the crossover on the energy distribution shown in Fig. 2. The sharp lower cutoff is due to the fixed activity threshold w_a .




Figure 2. Probability distributions of energy E , peak power W , and lifetime T of auroral emission events initiated poleward ($\phi > 0$, crosses) and equatorward ($\phi < 0$, stars) of the empirical IB. The low-latitude distributions are shifted for easier comparison. The solid lines show the log-log distribution slopes for small-scale LL events (the power-law portions to the left of the dashed vertical lines) and for the entire group of HL events.

Table 1. Comparative characteristics of low- and high-latitude emission events.

Type of events	LL ($\phi < 0$)	HL ($\phi > 0$)
# of events	1250 (16.7%)	6231 (82.3%)
E_{tot}, J	5.7×10^{15} (47.5%)	6.3×10^{15} (52.5%)
τ_E	1.96 ± 0.04	1.56 ± 0.02
τ_W	2.14 ± 0.07	1.80 ± 0.02
τ_T	3.30 ± 0.26	2.26 ± 0.13
τ_A	2.30 ± 0.10	1.93 ± 0.06

Table 2. Scaling exponent τ_E of emission events, solar wind dynamic pressure (P_d , auroral electrojet (AE) index and the magnetotail (MT) index (IB latitude at $MLT=0$, (eq. 1)), for four ranges of onset $MLat$.

$MLat$	τ_E	P_d , nPa	AE , nT	MT
< 65	2.06 ± 0.04	2.64 ± 0.05	206 ± 5	63.2 ± 0.03
$65...67$	1.88 ± 0.04	2.42 ± 0.03	106 ± 3	63.9 ± 0.03
$67...69$	1.59 ± 0.03	2.41 ± 0.03	84 ± 3	64.0 ± 0.03
> 69	1.58 ± 0.03	2.20 ± 0.03	93 ± 4	64.1 ± 0.03